

1993

REPRINTED FROM:

MARINE GEOLOGY

*International Journal of Marine Geology,
Geochemistry and Geophysics*

Marine Geology, 114 (1993) 155-170
Elsevier Science Publishers B.V., Amsterdam

Channels, echo character mapping and tectonics from 3.5 kHz profiles, distal Bengal Fan

Lisa J. Redbourn^a, Jonathan M. Bull^{a,1}, Roger A. Scrutton^b and Dorrik A.V. Stow^a

^a*Department of Geology, University of Southampton, Highfield, Southampton SO9 5NH, UK*

^b*Department of Geology and Geophysics, University of Edinburgh, West Mains Road, Edinburgh EH9 3JW, UK*

(Received August 20, 1992; revision accepted June 14, 1993)



ELSEVIER
AMSTERDAM - LONDON - NEW YORK - TOKYO

Editors

For Europe, Africa and the Near East:

H. Chamley, Université des Sciences et Techniques de Lille-Artois, Sédimentologie et Géochimie, B.P. 36, 59655 Villeneuve d'Ascq, France
Phone: 20 43 41 30

For the Americas, Pacific and Far East:

D.A. McManus, University of Washington, School of Oceanography WB-10, Seattle, WA 98195, U.S.A.

Editorial Board

F. Barbieri, Parma

W. Berger, La Jolla, Calif.

A.H. Bouma, Baton Rouge, La.

A.J. Bowen, Halifax, N.S.

L. Carter, Wellington

D.S. Cronan, London

J.R. Curray, La Jolla, Calif.

P.J. Davies, Sydney, N.S.W.

M.L. Delaney, Santa Cruz, Calif.

R.V. Dingle, Cape Town

S.L. Eitrem, Menlo Park, Calif.

M.E. Field, Menlo Park, Calif.

J. Francheteau, Paris

G.P. Glasby, Sheffield

D.S. Gorsline, Los Angeles, Calif.

M. Grant Gross, Baltimore, Md.

A. Guilcher, Brest

J.R. Hails, Wuhan

E.L. Hamilton, San Diego, Calif.

D.M. Hanes, Gainesville, Fla.

W.W. Hay, Boulder, Colo.

A.C. Hine, St. Petersburg, Fla.

H. Holteidahl, Bergen

D. Jongsma, O'Connor, A.C.T.

N.H. Kenyon, Godalming

E.M. Klein, Durham, N.C.

H.J. Knebel, Woods Hole, Mass.

I.N. McCave, Cambridge

R.E. McDuff, Seattle, Wash.

A.R.M. Nowell, Seattle, Wash.

R.Q. Oaks, Jr., Logan, Utah

O.H. Pilkey, Durham, N.C.

H. Postma, Texel

A.I. Rees, Godalming

H.E. Reineck, Wilhelmshaven

D.A. Ross, Woods Hole, Mass.

W.F. Ruddiman, Charlottesville, Va.

E. Seibold, Freiburg

J.M. Skey, Oslo

D.J. Stanley, Washington, D.C.

A.H.B. Stride, Godalming

D. Stüben, Kiel

D.J.P. Swift, Norfolk, Va.

S. Uyeda, Tokyo

J.T. Wells, Morehead City, N.C.

R.A. Wheatcroft, Woods Hole, Mass.

L.D. Wright, Gloucester Point, Va.

Channels, echo character mapping and tectonics from 3.5 kHz profiles, distal Bengal Fan

Lisa J. Redbourn^a, Jonathan M. Bull^{a,1}, Roger A. Scrutton^b and Dorrik A.V. Stow^a^aDepartment of Geology, University of Southampton, Highfield, Southampton SO9 5NH, UK^bDepartment of Geology and Geophysics, University of Edinburgh, West Mains Road, Edinburgh EH9 3JW, UK

(Received August 20, 1992; revision accepted June 14, 1993)

ABSTRACTRedbourn, L.J., Bull, J.M., Scrutton, R.A. and Stow, D.A.V., 1993. Channels, echo character mapping and tectonics from 3.5 kHz profiles, distal Bengal Fan. *Mar. Geol.*, 114: 155–170.

The distal parts of the Bengal Fan are spectacularly affected by tectonic deformation related to the diffuse plate boundary between the Indian and Australian plates. Here we use 3.5 kHz and 12 kHz echosounder profiles, seismic reflection profiles and piston core results to examine sedimentary processes and their relationships to tectonism within an area (78°–82°E, 0°–6°S) just to the south of Ocean Drilling Program Leg 116 sites.

Echo character mapping was completed using echosounder data in conjunction with results from piston coring, and a total of five different echo types have been recognised. Four of these fall into the echo character classification scheme developed by Damuth (1980a), whilst the fifth is believed to represent hemiturbidite deposits. Several types of submarine channel were also identified from echosounder data and a correlation between echo type and channel location can be seen. Their abundance, erosional and/or depositional nature together with a complex meandering and bifurcation pattern across a wide region of average gradient around 1°/km, are all features characteristic of a broad *channel termination zone* on a large elongate fan. It is clear, therefore, that the Bengal Fan extends beyond 6°S.

Active faulting in the area has led to the development of an irregular topography of low rounded hillocks that interfere with incoming turbidity currents. This has resulted in ponding between highs rather than lobe construction, thinning and pinching out of turbidites against the flanks of local relief, flow lofting and hemiturbidite drape, and common small-scale slumping. In some cases uplifted channel segments can be seen abandoned and partially filled.

Introduction

The Bengal Fan (Fig. 1) is one of the world's largest elongate submarine fans and has an area of 2.8–3 × 10⁶ km² (Curray and Moore, 1971; Curray et al., 1982; Emmel and Curray, 1984). Channels up to 2500 km long act as conduits for turbidity currents flowing the full length of the fan from its apex in the northern Indian Ocean off the Ganges Delta until it merges with the central abyssal plain south of the equator. The distal parts of this fan complex are spectacularly affected by deformation associated with the diffuse plate boundary zone across the northern Indian Ocean.

¹Author to whom all correspondence should be addressed

The objectives of this paper are to characterise the channel termination zone of the Bengal Fan and to study the complex interplay between sedimentation and tectonics in this unique area.

It is now recognised that the zone of diffuse deformation in the northern Indian Ocean, representing the plate boundary between the Indian and Australian plates, extends for thousands of kilometres from the Chagos Bank to the Sumatran Trench (Gordon et al., 1990). Within the Central Indian Ocean Basin this deformation manifests itself tectonically with an E–W fabric on two spatial scales: long wavelength (150–200 km) undulations of basement and overlying sediments; and faulted blocks 5–20 km in width which are bounded by high-angle reverse faults that extend throughout

Scope of the journal

Marine Geology is an international medium for the publication of original studies and comprehensive reviews in the field of Marine Geology, Geochemistry and Geophysics. The editors endeavour to maintain a high scientific level and it is hoped that with its international coverage the journal will contribute to the sound development of this field.

A *letter section* is provided as a publication outlet for short papers which require *rapid* publication.

Publication schedule and subscription information – 1993

Marine Geology (ISSN 0025–3227) is published as seven volumes a year. The subscription price for 1993 (Volumes 108–114) is Dfl. 2135 + Dfl. 231 p.h. = Dfl. 2366 (approx. US\$1315). The Dutch guilder price is definitive. The U.S. dollar price is subject to exchange-rate fluctuations and is given only as a guide. Subscriptions are accepted on a prepaid basis only, unless different terms have previously been agreed upon. Personal subscription rates and conditions are available upon request from the Publisher. Subscription orders can be entered only by calendar year (Jan.–Dec.) and should be sent to Elsevier Science Publishers, Journal Department, P.O. Box 211, 1000 AE Amsterdam, The Netherlands, tel. 31.20.5803642, fax 31.20.5803598, or to your usual subscription agent. Postage & handling (p.h.) charges include surface delivery, except to the following countries where air delivery via S.A.L. (Surface Air Lift) mail is ensured: Argentina, Australia, Brazil, Canada, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, PR China, Singapore, South Africa, South Korea, Taiwan, Thailand and the USA. For all other countries, airmail rates are available upon request. Claims for missing issues must be made within six months of publication (mailing) date; otherwise, such claims cannot be honoured free of charge. For further information, or a free sample copy of this or any other Elsevier Science Publishers journal, readers in the USA and Canada can contact the following address: Elsevier Science Publishing Co. Inc., Journal Information Center, 655 Avenue of the Americas, New York, NY 10010, U.S.A., tel. 212.63337502. Back issues can be ordered. For further information regarding prices etc., please contact Elsevier Science Publishers, Journal Department, P.O. Box 211, 1000 AE Amsterdam, The Netherlands, tel. 31.20.5803642, fax 31.20.5803598.

US mailing notice — *Marine Geology* (ISSN 0025–3227) is published monthly by Elsevier Science Publishers, (Molenwerf 1, Postbus 211, 1000 AE Amsterdam). Annual subscription price in the USA US\$1315 (subject to change), including air speed delivery. Second class postage paid at Jamaica, NY 11431.

USA POSTMASTERS: Send address changes to *Marine Geology*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003.

Airfreight and mailing in the USA by Publications Expediting.

(continued on inside back cover)

© 1993, ELSEVIER SCIENCE PUBLISHERS B.V., ALL RIGHTS RESERVED

0025-3227/93/\$06.00

No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Special regulations for readers in the U.S.A. — This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The per-copy fee is stated in the code-line at the bottom of the first page of each article. The appropriate fee, together with a copy of the first page of the article, should be forwarded to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, U.S.A. If no code-line appears, broad consent to copy has not been given and permission to copy must be obtained directly from the author(s). All articles published prior to 1980 may be copied for a per-copy fee of US \$2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the material herein.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such a product or of the claims made of it by its manufacturer.

This issue is printed on acid-free paper.

PRINTED IN THE NETHERLANDS

0025-3227/93/\$06.00

© 1993 — Elsevier Science Publishers B.V. All rights reserved.




Fig. 1. Location map showing position of survey area and ODP Leg 116 sites in the Central Indian Ocean. Bathymetry in metres (after Emmel and Curray, 1984).

the oceanic crust (Weissel et al., 1980; Bull, 1990; Bull and Scrutton, 1990, 1992).

The distal parts of the western Bengal Fan were the subject of a recent Ocean Drilling Program Leg (116) whose objectives were to investigate the timing and development of the deformation as well

as the sedimentary history of the Bengal Fan. The principal results of the Leg (Shipboard Scientific Party, 1989) are briefly discussed here.

The first of the objectives was met by drilling a pair of companion sites on one of the fault blocks in the deformation area. One of these sites, 717,

was in the thickest part of the axis of a syncline, whereas site 719 was partway up the block where syndeformational sediments are thinner. Sediments at both sites are fan turbidites with distinctive turbidite layers correlatable between the two sites. Reduction in thickness of the section between the two sites has occurred by pinching out of beds and thinning of individual turbidites. From the gradual nature of this reduction together with dating studies using calcareous nannofossils it was concluded that movement on the fault has been relatively slow and constant, perhaps with a slight increase with time. The onset of deformation was 7.5–8.0 Ma.

The second objective, the sedimentary history of the distal Bengal Fan, was met partly by sites 717 and 719 and partly by drilling on the next block south where the post-Miocene section is attenuated, thus allowing penetration of older sediments. At the base of this hole (site 718) fan sediments of early Miocene age (17 Ma) were still being penetrated with no evidence for reaching the base of the Fan. Almost the entire Miocene section consists of silt and silty mud turbidites.

The best studied area of the deformation zone is just to the south of the Leg 116 sites where there is now reasonable data coverage (see Fig. 2, for example). Authors have looked in detail at the tectonic deformation in this area from seismic reflection profiles (Neprochnov et al., 1988; Bull, 1990; Bull and Scrutton, 1990, 1992). These studies have concentrated on the deeper structure of the deformation. In this paper we use 3.5 kHz profiles, in combination with 12 kHz echograms, seismic reflection profiles and core data, to study sedimentary processes and their interaction with the two spatial scales of deformation. The dataset used in this study was collected by Lamont-Doherty Geological Observatory and Edinburgh University.

Seafloor bathymetry

Seafloor bathymetry is constrained by regularly spaced ship tracks in the south of the area (Fig. 2). In the north, the ship tracks are clustered around the ODP drilling sites, so the bathymetry is only well constrained in this small region. Seafloor

depths range from 4500 m in the north of the study area to 5100 m in the south, whilst just to the east the Afanasy-Nikitin Seamount rises some 2300 m above the surrounding basin. If the effect of short wavelength bathymetric variations is ignored, the depth increases from north to south across the area by 600 m over a distance of nearly 450 km, as would be expected at the distal end of a sedimentary fan. The most prominent short wavelength feature is a seafloor high, centred at 80°00'E and 04°15'S, which rises about 400 m above the surrounding seafloor. Its flanks slope at an angle of 0.6°. There is a smaller high, centred at 81°25'E and 04°45'S, which has a height of about 200 m and flanks which slope at 0.8°.

It is observed that the seafloor bathymetry closely follows the basement topography, as determined by Bull (1990), with seafloor undulations, where present, orientated E–W and apparently discontinuous across N–S orientated fracture zones.

Channel characteristics

In the surveyed area, 35 crossings of submarine channels have been identified from 3.5 kHz and 12 kHz echosounder and from seismic reflection sections (Fig. 3). The channels can be classified by their topographic relationship to the local mean surface of the fan. About 60% of the channels are incised below the local fan surface representing erosional or scoured features and the remaining 40% are built up above it representing depositional features (Normark, 1970). Channel depths range between 3 and 41 m, with 65% of channels having depths less than 10 m. As the angle of a ship's track across a channel is, in most cases, unknown, the channel widths calculated from echosounder profiles are unreliable. However, the maximum recorded apparent width of 3500 m places an upper limit on true channel widths. The minimum recorded width is 180 m. Around the ODP sites, the high density of ship tracks allows two erosional channel paths to be traced across a small distance (approximately 10 km). In this case, true widths of 750 m and 1090 m were calculated for the two channels. In general, the wide spacing of ship tracks relative to the presumed complexities of




Fig. 2. Map of bathymetry of survey area, and echo character of seafloor showing the distribution of the various echo types described in the text and shown in Fig. 5. Location of 3.5 kHz echograms (ship tracks) are shown.

channel pathways (Damuth et al., 1983; Carter, 1988), over most of the area prohibits accurate channel path determination. However, several other channel characteristics have been noted. The channel lettering employed below refers to Figs. 3 and 4.

Channel A is wide and shallow with levees which are only slightly raised. Over both levees, the echosounder profile shows a long acoustic return with a strong bottom echo and no sub-bottom reflections. The echo length decreases with increasing distance from the channel axis. In the channel




Fig. 3. Map showing positions of channel crossings on echosounder profiles, and areas of active faulting. The lettered channels are described in the text and shown in Fig. 4. Note that areas of active faults correlate with seafloor highs.

itself, the acoustic return is shorter although the bottom echo is still strong, and the lack of acoustic penetration and sub-bottom reflections suggest that the channel is floored by sandy sediments whose surface is broken up by bedforms produced by turbidity currents (Damuth, 1975). Active depositional channels are typically floored by sandy

sediments whose sand content decreases further away from the channel floor (Normark, 1978), and given the width and low relief of channel A, it is interpreted as an active depositional channel with levees which are just starting to develop.

Channel E is a good example of a more established depositional channel. Weak but coherent




Fig. 4. Diagram showing channel profiles recorded by 3.5 kHz echosounder with associated line drawings. (A) Active depositional channel, (B) erosional channel bounded to the right by a fault block, (C) infilled channel, (D) typical erosional or scoured channel, (E) typical depositional channel, (F) large depositional channel, thought to be one of the main distributary channels in survey area.

reflections are present in the levees and the channel itself and both levees are built up above the local mean surface of the fan.

In a small part of the area, centred at 80°50'E and 03°35'S, three large depositional channels appear on three different sections of the echosounder profile. Channel F is imaged by one of the sections. Some incoherent reflections are present in the levees and the channel floor comprises weak,

prolonged echoes. One of the channel walls appears to have slumped down to the channel floor, producing an asymmetric profile. On the basis of their similar seismic character, their equivalent dimensions and close physical proximity, these three imaged channels are almost certainly part of the same system, and this is by far the largest in the surveyed area with a depth of 41 m. Therefore, it may be one of the main active distributary channels

of the fan, in this area. The recorded positions of this channel indicate that it follows a meandering path, the wavelength of which appears to be 30 km over this small section.

Channel B is bounded to the right by a 10 m high fault block which comprises a strong seafloor echo with several continuous, parallel sub-bottoms, whereas weaker, prolonged echoes are returned from the channel floor itself. As the channel is flush with the base of the fault scarp, it seems likely that the channel path has been controlled by fault-induced bathymetry. Therefore, the faulting most likely occurred prior to the development of this part of the channel.

Within the surveyed area, there are several examples of channels which have been partly or completely infilled by younger sediments. In the case of channel C, the profile is visible as a weak sub-bottom reflection. The infilling sediments are acoustically transparent with a strong bottom echo. Damuth (1980a,b) suggested that this type of echogram is typical of a mass-wasting process, such as a debris flow. However, as the channel is situated at the very distal end of the fan where regional slopes are minimal, the debris flow was probably a local event derived from a fault block high, or the channel has been filled by homogeneous fine-grained muds.

The scoured, erosional channels mentioned earlier are typified by channel D. The echoes over this channel are weak and prolonged with a strong bottom return, indicating limited acoustic penetration. The channel appears "V"-shaped because of the vertical exaggeration on the echosounder profile. It has no levees and a faint diffraction pattern extends down from the base of the channel. In the surveyed area, this type of channel is no more than 10 m deep and usually has an apparent width of less than 1000 m. Thus, scoured channels appear to be narrower than depositional channels, whereas channel depths are similar for both types.

Most of the channels, including the main distributary one centred at 80°50'E and 03°35'S, are located in the deeper, flatter parts of the area. However, a partially infilled channel can be seen located at a point about 100 m below the summit of the main seafloor high already noted. In this channel, an acoustically transparent layer has been

deposited which onlaps onto the levees. The data quality along this section of 3.5 kHz echosounder profile is very poor, but several other channels have been tentatively identified over the summit of the seafloor high. These indistinct channels are assumed to be inactive as it is unlikely that active channels would continue up the inclined side of the seafloor high. A probable explanation for their occurrence is that they developed prior to the uplift of the seafloor high which is coincident with the crest of a long wavelength basement undulation. This uplift rendered them inactive and allowed them to become subsequently infilled by more recent sediments, presumably hemipelagic or hemiturbidite.

Two almost totally infilled channels have been identified in the north of the survey area near the ODP sites (one of them being channel C, which has been described earlier). In both cases, the old channel profile remains visible as a sub-bottom reflection and the infilling material is acoustically transparent. There are, at least, two major faults in this region, both of which downthrow towards the south. It is possible that the faults have undergone displacement in both a lateral, as well as vertical sense which may have diverted the paths of existing channels in the area, thus rendering some channel sections inactive.

Echo character mapping

Echo character mapping has been completed in the surveyed area using 3.5 kHz echosounder profiles. Five distinct echo types have been identified, and their nature and distribution throughout the area are shown in Figs. 2 and 5. The echo type interpretation follows Damuth (1975, 1980a,b).

Echo type 1 (Fig. 5) comprises a very prolonged echo with a strong seafloor return. No sub-bottom reflections are visible. This echo type appears to be the same as the class IIB echoes in the classification scheme developed by Damuth, which are associated with sediments comprising, on average, 54% silt/sand beds and 46% mud. It has been suggested that the prolonged echoes and lack of sub-bottoms, which characterize echo type 1, could be a result of scattering of acoustic energy either by erosional/depositional bedforms on the seafloor




Fig. 5. Diagram showing examples of the five echo types recorded by 3.5 kHz echosounder in the survey area, and described in the text. Type 1: very prolonged echo with a strong sea floor return and no sub-bottom reflections, type 2: prolonged echo with widely-spaced, coherent, parallel sub-bottom reflections, type 3: slightly prolonged echo with indistinct, discontinuous sub-bottom reflections, type 4: multiple distinct, closely-spaced, parallel sub-bottom reflections, type 5: acoustically transparent layer with weak sea floor echo. See text for further discussion.

and/or by relatively thick sand/silt beds (Damuth, 1975, 1978, 1980a; Damuth and Hayes, 1977; Embley, 1976). If the acoustic energy is scattered at the seafloor, then less energy is transmitted and the prolonged echo from scattering masks sub-bottom reflections. Damuth suggests that high velocity turbidity currents are responsible for producing the large amounts of silt/sand present in the sediments, and also for producing erosional/depositional bedforms. Thus, echo type 1 is interpreted as sediments of mainly turbiditic origin.

Echo type 2 (Fig. 5) comprises a prolonged echo with weak, widely spaced, coherent, parallel sub-bottom reflections. Although this echo type is not included in Damuth's classification scheme, it is most similar to the class IIB echoes; the differences being that echo type 2 has a slightly less prolonged echo with one or two sub-bottoms present. These differences are thought to indicate that there is less silt/sand present than in echo type 1 and also that acoustic energy is still being scattered at the seafloor, but to a lesser extent than in echo type 1, so the depth to which sub-bottoms are recorded increases. The reduction in seafloor scattering implies that there may be fewer erosional/depositional bedforms present and therefore, the sediments responsible for echo type 2 may have been deposited by weaker and/or more muddy turbidity currents than in the case of type 1 echoes.

Type 3 echoes (Fig. 5) comprise a slightly prolonged echo with indistinct, generally discontinuous sub-bottoms of varying lateral extent, and a clear sea floor reflection. This description fits the class IIA echoes in Damuth's classification scheme. The average bedded silt/sand content of sediments generating type 3 echoes was found to be 11% (Damuth, 1980a). The echo type is thought to be caused by the presence of erosional/depositional bedforms patchily distributed across the seafloor, which results in a differential scattering of acoustic energy and, therefore, semi-prolonged echoes with intermittent sub-bottoms are recorded.

Echo type 4 (Fig. 5) comprises multiple distinct, closely spaced, parallel sub-bottom reflections which are mostly laterally coherent, although they vary in reflectivity. There are hardly any prolonged echoes and the sea floor reflection is clear and continuous. This echo type falls into class IB of

Damuth's classification scheme, and the associated sediments have an average bedded silt/sand content of 3% (Damuth, 1980a). According to Damuth, echoes can be generated by interference from many thin layers and also be produced by factors such as changes in density and CaCO_3 content, and differential sediment compaction and grain size. Therefore, the sub-bottoms are not directly related to individual beds. The paucity of silt/sand beds suggests that sediments generating type 4 echoes are either less controlled by turbidity currents or, more likely, are entirely mud-rich turbidites. These are generally found in the more distal regions of the fan and adjacent to the main distributary channel pathways.

It can be seen from the previous descriptions that echo types 1 and 4 represent the end members of a continuous sequence. Progressing from type 1 to type 4, the echoes become less prolonged and sub-bottoms appear and become more closely-spaced and coherent. These echo variations correspond to a decrease in the bedded silt/sand content of the sediment and possibly to a reduction in the number of erosional/depositional bedforms present on the seafloor, amongst other factors. Thus, the sequence of echo types from 1 to 4, is interpreted to represent environments of deposition which are progressively more mud rich. Turbidity currents have progressively lost their silt/sand load and mud turbidites are more interbedded with thick hemipelagic and hemiturbidite deposits. They are generally more distal in setting.

Echo type 5 comprises a weak sea floor echo and is internally acoustically transparent. Along some parts of the echosounder profile, the acoustically transparent layer overlies sediments with a different echo character. Its thickness can be up to 20 m but is usually less than 10 m. The transparent layer occurs over a small region which is centred at $79^{\circ}30'E$ and $04^{\circ}15'S$ and lies immediately west of the main sea floor high (Fig. 2). The transparent layer has been recorded in this region by both the *Vema* 2902 and *Robert Conrad* 2706 cruises.

Along two sections of RC2706 echosounder profile, both of which are orientated approximately N-S, the transparent layer has accumulated in the form of a drape deposit, up the north-facing slopes of fault blocks and small topographic highs only.

This suggests that the acoustically transparent sediments were deposited from a southward travelling flow. However, preferentially accumulated transparent sediments are not seen on any other sections of profile over the region, so it is possible that the southward flow was very weak, or that localized reworking of sediments by bottom currents is responsible for the observed accumulations.

It is suggested that echo type 5 could be generated by a series of hemiturbidite deposits, as described by Stow and Wetzel (1990). They define a hemiturbidite as a very fine-grained muddy sediment with no distinct bedding, partly turbiditic, partly hemipelagic in character that is deposited from an essentially stationary, dilute suspension cloud which is formed from, but beyond and above the dying stages of a large turbidity current. Hemiturbidites are most likely to occur in distal turbidite settings and can be deposited over a period of at least a few months for a unit 1 m thick. Consequently they are bioturbated throughout most of their thickness and this bioturbation causes complete homogenisation of these very fine-grained sediments rendering them acoustically transparent internally.

Seismic p-wave velocities were found to be anisotropic at the base of some hemiturbidite beds (Stow and Wetzel, 1990) which indicates that the lowermost units of these beds were deposited from a very dilute turbidity current. The same dilute nature could explain the accumulations of acoustically transparent sediments up north-facing slopes that originated from generally southward directed flows, and also the acoustically transparent fills of some abandoned channels. However, the confinement of the sediment fill to the channel axes argues against a true hemiturbidite deposit in this case.

Sediment cores and their relationship to echo character

Within the surveyed area, the results from three Ocean Drilling Program sites and eleven piston cores taken by the Lamont-Doherty Geological Observatory were available. The sediments recovered from all fourteen sites are dominated by

turbiditic inputs with only very minor pelagic intercalation.

The boreholes from the three closely spaced ODP sites (717, 718 and 719) drilled at approximately 81°24'E and 01°00'S, all show very comparable lithostratigraphic sections (Stow et al., 1990). These are dominated by turbidites from four main sources: thin to very thick-bedded silt/sand and mud turbidites derived from a Himalayan source; thin to thick dark mud turbidites derived from the east Indian continental margin having a marked Deccan Traps signature; nannofossil mud turbidites from the south Indian and Sri Lankan margin; and pale foraminiferal-rich turbidites from a local seamount source. Pelagic intercalations are minimal but thick sections of hemiturbidite deposits occur in parts. This facies association extends to the base of each borehole drilled, i.e. to a depth below seafloor of 961 m in the longest borehole.

The eleven piston cores have a length which ranges between 48 cm and 964 cm and has an average value of 472 cm. The core sites are widely dispersed over the surveyed area and the recovered sediments can be broadly divided into three categories (see Table 1). One core in the north of the area (RC17130) comprises layers of foraminiferal marl, and interbedded terrigenous clay and foraminiferal marl. These sediments appear to be the product of alternating pelagic and turbiditic inputs.

The second category consists of cores with sandier sediments. Cores VM29037 and VM19173 are situated approximately in the middle of the area and comprise sediments with nearly 100% quartz sand content overlain by yellow/brown coloured clay, which is approximately 0.7 m thick, with less than 1% sand content. Further to the south, core VM29038 comprises sediments with a slightly lower quartz sand content (between 70% and 95%), present throughout the core. Also in the south of the area, core VM19172 consists of interbedded silt and sand/silt zones which have sand contents of 30% and 85%, respectively. Sediments generally have a lower coarse fraction as they are deposited progressively further away from their terrigenous source, with the exception of sediments from core RC17130, and this is consistent with the observed decrease in sand content from the centre of the area southwards.

TABLE 1

Piston core information

Piston core	Location	Description
RC17130	80°09'E, 00°56'S	Foraminiferal marl with some interbedded clay
VM19172	80°30'E, 05°32'S	Interbedded silt and sand with manganese nodules
VM19173	80°35'E, 03°15'S	Grey quartz sand
VM29010	80°52'E, 01°19'S	Green/grey fine-grained clay with manganese nodules
VM29037	79°11'E, 03°00'S	Grey clay overlying dark grey sand
VM29038	78°39'E, 05°43'S	Green/grey sand with manganese nodules
VM33055	81°42'E, 04°44'S	Grey/brown radiolarian clay
VM33057	81°41'E, 01°30'S	Yellow/grey clay, laminated in places
VM36057	81°38'E, 02°34'S	Yellow/brown clay with manganese nodules
VM36058	81°37'E, 02°31'S	Brown clay with manganese nodules and volcanic mud
VM36059	81°37'E, 02°29'S	Brown clay with manganese nodules and volcanic mud

The main distributary channel identified in the area is located near to one of the cores with a high sand content.

The remainder of the piston cores consist of clays which have a sand content of between 0% and 3% and are rarely laminated. The clays contain varying proportions of radiolaria, volcanic glass shards, sponge spicules, mica and quartz. In two of these cores, VM36058 and VM36059 situated in the east of the area near the base of the Afanasy-Nikitin seamount, an 18 cm thick layer of volcanic mud is present at the top of the cores which could be the result of a recent seamount eruption, possibly resedimented down the side of the seamount. However, the rest of the sediments in this category are mostly terrigenous turbidites, with predominant micaceous silts and muds that originated from the Himalayas and reached the Bengal Fan via the Ganges Delta (Curry and Moore, 1971; Stow and Cochran, 1989).

Out of the eleven piston cores, seven contained manganese micronodules, although there does not

appear to be any pattern in the distribution of these cores over the surveyed area. The presence of micronodules is consistent with the findings of Bannerjee and Mukhopadhyay (1991) who observed that small (<4 cm diameter) manganese nodules are most common north of 9°S, which would include this study area.

Faulting

The recent faults in the survey area, which have been interpreted from 3.5 kHz and 12 kHz echosounder data, and seismic reflection profiles, fall into two categories. The majority of faults are expressed at the surface by rounded "hillocks" in the seafloor bathymetry (Fig. 6). In general, the base of one side of the hillock appears to be almost continuous with the surrounding seafloor, whereas, on the other side of the hillock, its base is delineated by a diffraction pattern which extends below the seafloor. Seismic reflection profiles (Fig. 6, see Bull and Scrutton, 1992, for more discussion on their tectonic significance) show that these hillocks are, in fact, the hanging wall anticlines of the faults, with the diffraction pattern on the faulted side. In most cases, little or no structure can be seen in the hanging wall anticlines on echosounder profiles, but some display several small, closely spaced faults over their surface.

The major faults, which produce the hanging wall anticlines, have been seen on seismic reflection profiles to extend throughout the sedimentary cover and down into the oceanic crust (Bull and Scrutton, 1990, 1992). Echosounder profiles show that echoes from over the anticlines are usually weaker and shorter than echoes from the surrounding seafloor. This is possibly a result of slightly finer sediments being deposited post-faulting on these relative highs. Some faults only appear as an offset in seafloor depth on echosounder records, but coincident reflection profiles demonstrate that they are real and also extend down to the basement. Basement faults which have not propagated upwards through the overlying sediment are imaged only by seismic reflection profiles and therefore, in areas surveyed by echosounder only, undetected basement faults are certainly present.

The maximum displacement of seafloor on either

side of the fault, or fault block, is found to be 130 m in this area, with an average of about 30 m. The widths of the fault blocks range between 1 km and 16 km with an average of about 4 km, and the fault blocks are tilted up to 5° above horizontal, the average value being almost 2°. The downthrow side of each fault is taken to be the side on which the seafloor is deepest. Within the survey area, 65% of all faults were found to downthrow towards the south, with the remainder downthrowing in a northerly direction. Bull (1990) found, on the basis of a seismic reflection profile study, that approximately equal numbers of faults downthrow south as downthrow north in this area. The discrepancy in these observations may be a product of the increased echosounder coverage in this study affording a more representative sample of the faults. Alternatively, it may indicate that faults downthrowing to the south have propagated further through the sedimentary cover and are therefore the most recently active and are the ones recognised preferentially on high-resolution (3.5 kHz) profiles.

The level of recent activity of the faults has been roughly assessed following the criteria that the more recent faults are sharply defined on the echosounder profiles, with little or no erosion over the fault block surface and no build up of sediments against the sides of the fault block. Three regions in which there are high numbers of active faults have been identified at 81°30'E, 02°50'S, 80°00'E, 04°15'S and 81°25'E, 04°45'S (see Fig. 3). The latter two regions are situated on the two seafloor highs described earlier, which correlate with basement highs. The other region is situated on the south facing flank of an E-W trending undulation in the seafloor.

The number of faults, both active and inactive, is markedly higher over these three regions and in general, there is a correlation between higher fault densities and the shallower parts of the survey area. Although this correlation is observed over the peaks of the E-W trending seafloor undulations in the west of the area, to some extent it may be a result of recent sediment infilling the troughs of the undulations, and therefore obscuring some of the faulting within these troughs. This is shown to be the case by seismic reflection




Fig. 6. One of the spectacular reverse faults within the intraplate deformation zone (at $81^{\circ}27'E$, $02^{\circ}29'S$) as revealed on (A) 3.5 kHz echosounder record, and (B) a migrated multichannel (12-fold) seismic reflection profile. The fault plane has a dip of 35° – 40° within oceanic basement (the top of which is denoted by the strong reflector at ca. 7.4 s TWT). The dip steepens above basement and there are a complex set of faults within the sedimentary cover. The 3.5 kHz record shows evidence of large-scale slumping. See Bull and Scrutton (1990, 1992) for more discussion of the seismic reflection profiles.

profiles, on which basement faults are imaged within the troughs. These faults have not affected overlying sediments and are not seen on echosounder records. However, overall the correlation between fault activity and seafloor/basement highs is believed to be real.

Discussion

Although the region surveyed in detail in this study is like a postage-stamp area at the end of the world's largest submarine fan, it nevertheless covers an area of 150,000–200,000 km². This is larger than most known modern fans and certainly far larger than almost all ancient turbidite successions that have been interpreted as submarine fans (Bouma et al., 1985a). It is important to bear in mind these scaling factors when considering the implications of this research to our understanding of fans in general (Bouma et al., 1985b). There are two aspects we wish to discuss further: (1) the nature of a channel-termination zone on a large mud-rich, elongate fan, and (2) the influence of within fan neotectonics on sedimentation.

Channel termination zone

According to previous work (Emmel and Curray, 1984; Stow and Cochran, 1989), the study area was beyond the reach of active channels at the feather edge of the fan where it merges with the Central Indian Ocean abyssal plain, over 2500 km from its apex in the north off the Ganges delta. However, this work has shown that the fan channel system extends further south than previously recognised and that the region studied is, in fact, one broad zone of channel termination.

The chief characteristics of this zone are illustrated in Fig. 7 and include the following:

(a) There is a very low average down-fan gradient (< 1 m/km), with a locally irregular topography related to channel margins and structural highs (see below).

(b) There are a large number of low-relief channels, mostly < 10 m deep but ranging up to 40 m in depth, with widths varying from < 200 m to > 1500 m. A great variety of channel types is evident, including broad flat-bottomed, narrow

V-shaped, sediment-filled or unfilled, erosional and/or constructional. There is a marked lack of direct continuity of channels from one survey crossing to another that suggests meandering, bifurcation and terminations are common.

(c) Channel levees built up above the surrounding fan surface, by between 2 m and 25 m, are found flanking just less than half the channels surveyed and are typically a few kilometres in width. In all cases levee heights are equivalent on left and right banks, attesting to the absence of a strong Coriolis force influencing deposition at these equatorial latitudes.

(d) The echocharacter distribution is complex. Four of the echocharacter types fall into the classification scheme developed by Damuth (1980a) and are readily interpreted in terms of turbidite deposition, ranging from more sand/silt-rich and higher energy currents (Type 1) to more mud-rich and lower energy currents (Type 4). There is a general N–S trend from Types 1–2 to Types 3–4, and also a topographic control with Types 1–2 occupying the lower ground. Both of these observations are consistent with a decrease in the influence and/or competence of turbidity currents across the channel termination zone, and with the fact that turbidity currents preferentially flow along bathymetric lows. Type 5 echoes are discussed below.

(e) The sediments recovered from the three ODP sites in the north of the area and from the scattered piston cores to the south are dominantly turbiditic (Stow et al., 1990), including thin to very thick-bedded silt/sand and mud turbidites as well as nanofossil–foraminiferal bioclastic turbidites. Hemiturbidites were common in parts of the ODP boreholes (Stow and Wetzel, 1990) but pelagic sediments extremely rare. Turbidites from at least three distinct sources have been recognised at the ODP sites, showing that whatever is fed in the upper reaches of the fan system finds its way through to the channel termination zone.

Although the data are sparse regionally, there is a general southward decrease in sand/silt content observed in the cores that is consistent with the downfan change in echocharacter noted previously.

All the features noted above (a–e) might be




Fig. 7. Schematic diagram of the Channel Termination Zone showing inferred channel pathways (based on actual channel crossings and echocharacter mapping), areas of positive relief related to tectonic activity, the Afanasy-Nikitin Seamount, and the nature of sedimentation resulting from interaction between current input and tectonic relief. Bathymetry of the Afanasy-Nikitin Seamount adapted from Sclater and Fisher (1974) using CD28 bathymetric data.

considered as typical of a channel termination zone from any submarine fan, and particularly those of large, muddy elongate fans (Stow, 1981, 1986). Similar characteristics have been described from the Mississippi (O'Connell et al., 1991), Amazon (Damuth et al., 1983), Indus (Kolla and Coumes, 1987) and Laurentian (Piper et al., 1983) fans, as well as from a number of smaller fan systems (Normark, 1978; Bouma et al., 1985a; Droz and Bellaiche, 1985). The chief characteristic is one of complexity, in morphology, echocharacter and sediments, but with an organisation that is logical both spatially and temporally. Channel termination on smaller (radial) fans is most commonly in the middle fan region and is characterised by rapid dumping of sand-rich sediments, whereas

on larger (elongate) fans the zone is more extended and the sediments finer grained.

However, there are several characteristics of the study area that we believe demonstrate the influence of within-fan neotectonic control on the otherwise normal pattern of deposition in a channel termination zone.

Neotectonics and sedimentation

The distal parts of the Bengal Fan lie within a broad area of active intraplate deformation that involves both the oceanic crust and overlying sediments (Bull and Scrutton, 1990, 1992). This has resulted in the otherwise "smooth" low-gradient fan surface being interrupted by a series

of rounded hillocks that are the surface expression of recent faulting. These create a local relief of typically 100–300 m with gradients up to 5° above the horizontal. The interaction of this irregular topography with incoming turbidity currents results in several unique effects.

(a) Channels are diverted around the base of upstanding hillocks, and this may help contribute to channel meandering, switching and abandonment, although such processes also occur under normal conditions (e.g. Kolla and Coumes, 1987). Channel remnants are left abandoned and now filled or partially filled on the flanks and crests of the hillocks, most likely as the direct result of tectonic uplift. The relative abundance of small erosional channels may be the result of local increase in flow velocities due to flow confinement around and/or over some of the hillocks.

(b) The complete lack of any detectable lobes in the whole of this channel termination zone is unusual for such a setting (e.g. Normark et al., 1983) and we suggest that this results from topographic interference to incoming flows. Deposition is concentrated in the low-lying areas and is more ponded than lobate, with individual turbidites thinning up onto and pinching out against adjacent hillocks.

(c) The thinning and pinching out of turbidites over highs has been carefully documented in the region of the ODP sites (Stow et al., 1990), as also has the occurrence of hemiturbidites (Stow and Wetzel, 1990). Although flow lofting and upward mixing of fine-grained sediment into the overlying water column is most likely a normal process for far-travelled turbidity currents (Sparks et al., 1993), the distal Bengal Fan is the first area from which the resulting hemiturbidite deposits have been described. We suggest that the presence of irregular hillocks in this area serves to enhance the flow lofting process.

Echotype 5 is interpreted as reflecting an accumulation of hemiturbidites that have been deposited on the western flank of the main seafloor high as a result of flow lofting from southerly flowing westward diverted dilute turbidity currents. This inference remains circumstantial until corroborated by further coring in the area.

(d) There seems to be an abundance of small-

scale slumping in the area as evidenced by chaotic reflectors associated with the margins of channels and with the more prominent hillocks. This local mass movement will be enhanced both by the presence of marked relief and by seismic activity related to fault displacement.

Although none of these features (a–d) are exclusive to zones of within-fan neotectonics, their relative abundance and co-occurrence are believed to be significant. The added influence of local relief and seismic activity to a channel termination zone will make an already highly varied depositional system still more complex in nature.

Acknowledgements

The authors are grateful to Jeff Weissel and Randy Lotti at Lamont-Doherty Geological Observatory for providing access to 3.5 kHz records and core data. The comments of two anonymous reviewers helped to greatly improve the manuscript. Edinburgh University data was collected during *Charles Darwin* cruise 28 and funded by NERC grant GR3/6480. LJR and this study were supported by NERC grant GR9/441.

References

- Bannerjee, R. and Mukhopadhyay, R., 1991. Nature and distribution of manganese nodules from three sediment domains of the Central Indian Basin, Indian Ocean. *Geo-Mar. Lett.*, 11: 39–43.
- Bouma, A.H., Normark, W.R. and Barnes, N.E., 1985a. *Submarine Fans and Related Turbiditic Systems*. Springer, New York, 351 pp.
- Bouma, A.H., Normark, W.R. and Barnes, N.E., 1985b. COMFAN: Needs and Initial Results. In: A.H. Bouma, W.R. Normark and N.E. Barnes (Editors), *Submarine Fans and Related Turbiditic Systems*. Springer, New York, pp. 7–12.
- Bull, J.M., 1990. Structural style of intra-plate deformation, Central Indian Ocean Basin: evidence for the role of fracture zones. *Tectonophysics*, 184: 213–228.
- Bull, J.M. and Scrutton, R.A., 1990. Fault reactivation in the Central Indian Ocean and the rheology of oceanic lithosphere. *Nature*, 344: 855–858.
- Bull, J.M. and Scrutton, R.A., 1992. Seismic reflection images of intraplate deformation, Central Indian Ocean basin, and their tectonic significance. *J. Geol. Soc. London*, 149: 955–966.
- Carter, R.M., 1988. The nature and evolution of deep-sea channel systems. *Basin Res.*, 1: 41–54.

- Curry, J.R. and Moore, D.G., 1971. Growth of the Bengal deep-sea fan and denudation in the Himalayas. *Geol. Soc. Am. Bull.*, 82: 563-572.
- Curry, J.R., Emmel, F.J., Moore, D.G. and Raitt, R.W., 1982. Structure, tectonics and geological history of the North Eastern Indian Ocean. In: A.E.M. Nairn and F.G. Stehli (Editors), *The Indian Ocean. (The Ocean Basins and Margins, 6.)* Plenum Press, New York, pp. 399-450.
- Damuth, J.E., 1975. Echo character of the Western equatorial Atlantic floor and its relationship to the dispersal and distribution of terrigenous sediments. *Mar. Geol.*, 18: 17-45.
- Damuth, J.E., 1978. Echo character of the Norwegian-Greenland sea: relationship to Quaternary sedimentation. *Mar. Geol.*, 28: 1-36.
- Damuth, J.E., 1980a. Use of high frequency (3.5-12 kHz) echograms in the study of near-bottom sedimentation processes in the deep-sea: a review. *Mar. Geol.*, 38: 51-75.
- Damuth, J.E., 1980b. Quaternary sedimentation processes in the South China Basin as revealed by echo character mapping and piston-core studies. In: D.E. Hayes (Editor), *The Tectonic/Geologic Evolution of Southeast Asian Seas and Islands*. Am. Geophys. Union Monogr., 23: 105-125.
- Damuth, J.E. and Hayes, D.E., 1977. Echo character of the East Brazilian continental margin and its relationship to sedimentary processes. *Mar. Geol.*, 24: 73-95.
- Damuth, J.E., Kolla, V., Flood, R.D., Kowsmann, R.O., Monteiro, M.C., Gorini, M.A., Palma, J.J.C. and Belderson, R.H., 1983. Distributary channel meandering and bifurcation patterns on the Amazon deep-sea fan as revealed by long-range side-scan sonar (GLORIA). *Geology*, 11: 94-98.
- Droz, L. and Bellaiche, G., 1985. Rhone deep-sea fan: morpho-structure and growth pattern. *AAPG Bull.*, 69: 460-479.
- Embley, R.W., 1976. New evidence for the occurrence of debris-flow deposits in the deep sea. *Deep-Sea Res.*, 17: 267-270.
- Emmel, F.J. and Curry, J.R., 1984. The Bengal submarine fan, Northeastern Indian Ocean. *Geo-Mar. Lett.*, 3: 119-124.
- Gordon, R.G., DeMets, L. and Argus, D.F., 1990. Present day motion between the Australian and Indian plates: kinematic constraints on distributed lithospheric deformation in the Equatorial Indian Ocean. *Tectonics*, 9: 409-423.
- Kolla, V. and Coumes, F., 1987. Morphology, internal structure, seismic stratigraphy and sedimentation of Indus fan. *AAPG Bull.*, 71: 650-677.
- Kolla, V., Moore, D.G. and Curry, J.R., 1976a. Recent bottom-current activity in the deep Western Bay of Bengal. *Mar. Geol.*, 21: 255-270.
- Kolla, V., Sullivan, L., Streeter, S.S. and Langseth, M.G., 1976b. Spreading of Antarctic Bottom Water and its effects on the floor of the Indian Ocean inferred from bottom-water potential temperature, turbidity and sea-floor photography. *Mar. Geol.*, 21: 121-189.
- Neprochnov, Y.P., Levchenko, O.V., Merklin, L.R. and Sedov, V.V., 1988. The structure and tectonics of the intraplate deformation area in the Indian Ocean. *Tectonophysics*, 156: 89-106.
- Normark, W.R., 1970. Growth patterns of deep-sea fans. *AAPG Bull.*, 54: 2170-2195.
- Normark, W.R., 1978. Fan valleys, channels and depositional lobes on modern submarine fans: characters for recognition of sandy turbidite environments. *AAPG Bull.*, 62: 912-931.
- Normark, W.R., Gutmacher, C.E., Chase, T.E. and Wikle, P., 1983. Monterey Fan: Growth pattern control by basin morphology and changing sealevels. *Geo-Mar. Lett.*, 3: 93-100.
- Piper, D.J.W., Normark, W.R. and Stow, D.A.V., 1983. The Laurentian Fan-Sohm Abyssal Plain. *Geo-Mar. Lett.*, 3: 141-146.
- O'Connell, S., Ryan, W.B.F. and Normark, W.R., 1991. Evolution of a fan channel on the surface of the outer Mississippi Fan: evidence from side-looking sonar. In: *Seismic Facies and Sedimentary Processes of Submarine Fans and Turbidite Systems*. Springer, New York, pp. 365-381.
- Sclater, J.G. and Fisher, R.L., 1974. Evolution of the east Central Indian Ocean, with emphasis on the tectonic setting of the Ninetyeast Ridge. *Geol. Soc. Am. Bull.*, 85: 683-702.
- Shipboard Scientific Party, 1989. ODP Leg 116 site survey. *Proc. ODP, Init. Rep.*, 116: 197-210.
- Sparks, R.S.J., Bonnezaze, R.T., Huppert, H.E., Lister, J.R., Hallworth, M.A., Mader, H. and Phillips, J., 1993. Sediment-laden gravity currents with reversing buoyancy. *Earth Planet. Sci. Lett.*, 114: 243-257.
- Stow, D.A.V., 1981. Laurentian Fan: morphology, sediments, processes and growth pattern. *AAPG Bull.*, 65: 375-393.
- Stow, D.A.V., 1986. Deep clastic seas. In: H.G. Reading (Editor), *Sedimentary Environments and Facies*. Blackwell, Oxford, 2nd ed., pp. 399-444.
- Stow, D.A.V. and Wetzel, A., 1990. Hemiturbidite: a new type of deep water sediment. *Proc. ODP, Sci. Results*, 116: 25-34.
- Stow, D.A.V., Cochran, J.R. and ODP Leg 116 Shipboard Scientific Party, 1989. The Bengal Fan: some preliminary results from ODP drilling. *Geo-Mar. Lett.*, 9: 1-10.
- Stow, D.A.V., Amano, K., Balson, P.S., Brass, G.W., Corrigan, J., Raman, C.V., Tiercelin, J.-J., Townsend, M. and Wijayananda, N.P., 1990. Sediment facies and process on the distal Bengal Fan, Leg 116. *Proc. ODP, Sci. Rep.*, 116: 377-396.
- Weissel, J.K., Anderson, R.N. and Geller, C.A., 1980. Deformation of the Indo-Australian plate. *Nature*, 287: 284-291.

(continued from inside front cover)

Publication of abstracts and contents lists

Abstracts and/or contents lists of this journal are published in the following journals: *Geoscience Contents*, *Geo Abstracts*, *Current Contents*, *Contents*, *Bulletin Signalétique*, *Chemical Abstracts*, *Marine Science Contents Tables*, *A.G.I.'s Bibliography and Index of Geology*, *Aquatic Sciences and Fisheries Abstracts*, *PASCAL/CNRS*, *Curr. Awareness Biol. Sci. (CABS)* and *Oceanic Abstracts*.

Note to contributors

A detailed *Guide for Authors* is available upon your request, and will also be printed in the first volume to appear each year. You are kindly asked to consult this guide. Please pay special attention to the following notes:

Language

The official language of the journal is English, but occasional articles in French and German will be considered for publication. Such articles should start with an abstract in English, headed by an English translation of the title. An abstract in the language of the paper should follow the English abstract. English translations of the figure captions should also be given.

Preparation of the text

- The manuscript should be typewritten with double spacing and wide margins and include at the beginning of the paper an abstract of not more than 500 words. Words to be printed in italics should be underlined. The metric system should be used throughout.
- The title page should include the name(s) of the author(s) and their affiliations.

Abstract

The abstract must be an informative statement of the content of the paper, explaining what the problem is, the methods used (when appropriate), a statement of the results, and the main conclusions. Constructions using phrases such as "this paper discusses", "are described" and "is reported" have no place in an abstract; the reader should be able to understand the essential qualities of the paper without referring to the paper itself. See K.K. Landes (*Bull. Am. Assoc. Pet. Geol.*, 1951, Vol. 35, p. 1660) for a concise statement on how to write a good abstract.

References

- References in the text start with the name of the author(s), followed by the publication date in parentheses.
- The reference list should be in alphabetical order and on sheets separate from the text.

Tables

Tables should be compiled on separate sheets. A title should be provided for each table and they should be referred to in the text.

Illustrations

- All illustrations should be numbered consecutively and referred to in the text.
- Drawings should be completely lettered, the size of the lettering being appropriate to that of the drawings, but taking into account the possible need for reduction in size (preferably not more than 50%). The page format of *Marine Geology* should be considered in designing the drawings.
- Photographs must be of good quality, printed on glossy paper.
- Figure captions should be supplied on a separate sheet.
- If contributors wish to have their original figures returned this should be requested in proof stage at the latest.
- Colour figures cannot normally be accepted unless the reproduction costs are to be met by the author. As a rough guide, the reproduction costs for a one-page figure are ca. Dfl. 2150 (inc. tax). Please consult the publisher for further information.

Proofs

One set of proofs will be sent to the author, to be checked for printer's errors. In case of two or more authors please indicate to whom the proofs should be sent.

Reprints

Fifty reprints of each article published are supplied free of charge. Additional reprints can be ordered on a reprint order form, which will be sent to the first author upon receipt of the accepted article by the publisher.

Submission of manuscripts

- Manuscripts originating in the Americas and Far East should be submitted in triplicate to Dr. D.A. McManus, Department of Oceanography, University of Washington, Seattle, Washington 98105, U.S.A. All other manuscripts should be sent in triplicate to the Editorial Office *Marine Geology*, P.O. Box 1930, 1000 BX Amsterdam, The Netherlands.
- Illustrations should also be submitted in triplicate. One set should be in a form ready for reproduction: the other two may be of lower quality.
- Authors are requested to submit the names and addresses of four potential referees with their manuscripts.
- Letter-type contributions should be submitted in triplicate to the Editorial Office *Marine Geology, Letter Section*, P.O. Box 1930, 1000 BX Amsterdam, The Netherlands. Contributions for this section should not exceed 8 printed pages and should be complete in themselves. No foldout illustrations can be accepted. The manuscripts received will be published ca. *three months* after acceptance. In order to achieve rapid publication, no proofs will be sent to the authors. Manuscripts should therefore be prepared with the greatest possible care.
- The indication of a FAX number on submission of the manuscript could assist in speeding communications. The FAX number for the Amsterdam office is: 020-5862696.
- In order to enable publication as quickly as possible after acceptance, authors are encouraged to submit the final text also on a 3.5" or 5.25" diskette. Both double density (DD) and high density (HD) diskettes are acceptable. Make sure, however, that the diskettes are formatted according to their capacity (HD or DD) before copying the files onto them. As with the requirements for manuscript submission, the main text, list of references, tables and figure captions should be stored in separate text files with clearly identifiable file names. The format of these files depends on the wordprocessor used. Texts written with DisplayWrite, MultiMate, Microsoft Word, Samna Word, Sprint, Volkswriter, Wang PC, WordMARC, WordPerfect and Wordstar or supplied in DCA/RFT or DEC/DX format can be readily processed. In all other cases the preferred format is DOS text or ASCII. Essential is that the name and version of the wordprocessing program, the type of computer on which the text was prepared, and the format of the text files are clearly indicated. **Authors are requested to ensure that the contents of the diskette correspond exactly to the contents of the hardcopy manuscript. Discrepancies can lead to proofs of the wrong version being made.**

Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere.

Upon acceptance of an article by the journal the author(s) will be asked to transfer the copyright of the article to the publishers. This transfer will ensure the widest possible dissemination of information under the U.S. Copyright Law.